

Crystal and molecular structure of (S)- α -(pCyanobenzenesulphonamido)- β -propiothiolactone, C₁₀H₈N₂O₃S₂

Matijašić, Ivanka; Andreotti, Giovanni D.; Sgarabotto, Paolo; Bezjak, Aleksandar; Fleš, Dragutin

Source / Izvornik: *Croatica Chemica Acta*, 1987, 60, 285 - 291

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:163:332207>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-10-07**



Repository / Repozitorij:

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



CCA-1726

YU ISSN 0011-1643

UDC 548.734.842.4

Original Scientific Paper

**Crystal and Molecular Structure of
(S)- α -(*p*-Cyanobenzenesulphonamido)- β -propiothiolactone,
 $C_{10}H_8N_2O_3S_2$**

Ivanka Matijašić

*Laboratory of Organic Chemistry and Biochemistry, Faculty of Science,
University of Zagreb, Strossmayerov trg 14, Zagreb,*

Giovanni D. Andreetti and Paolo Sgarabotto

*Istituto di Strutturistica Chimica, Università di Parma, C. N. R., Centro di Studio
per la Strutturistica Diffrattometrica, Parma, Italy,*

Aleksandar Bezjak

*Department of Chemistry, Faculty of Pharmacy and Biochemistry,
University of Zagreb, Zagreb,*

and

Dragutin Fleš

Research and Development Institute INA, Zagreb, Yugoslavia

Received July 7, 1986

Crystals of the title compound are monoclinic, space group $P2_1$, with $a = 1.0018(2)$, $b = 0.9722(2)$, $c = 0.6262(1)$ nm, $\beta = 102.32(1)^\circ$ and $Z = 2$. The structure was solved by direct methods using 1113 reflections with $I \geq 2\sigma(I)$ and refined to a conventional R index of 0.051. The heterocyclic four-membered ring is bent and has a puckering angle of $18.7(7)^\circ$. The dihedral angle between the best planes of the β -propiothiolactone and benzene ring is $70.2(3)^\circ$. Discrete molecules are connected along the b axis by intermolecular hydrogen bonds $N-H \dots O$ of 309.5(9) pm.

INTRODUCTION

The structure of the title compound was determined as a part of a continuing study of the α -substituted β -propiothiolactones.¹⁻⁴ These optically active compounds polymerized in solution and in bulk giving the polymers⁵⁻⁷ which represent a new class of nonamide-bonded polycysteins. The rate of polymerization is dependent on the nature of the substituent in the p -position on the benzene ring^{8,9}, which is, however, quite far from the atoms involved in polymerization.

This X-ray study was undertaken in order to establish the possible effect of the p -substituent on the geometry of β -propiothiolactone ring and of intermolecular contacts on bulk polymerization¹⁰.

EXPERIMENTAL

The crystals are colourless prisms elongated along [010]. Preliminary cell dimensions and space group data were obtained from oscillation and Weissenberg photographs. Lattice parameters were refined by least-squares using 15 reflections taken on a Siemens AED single-crystal diffractometer.

Crystal Data. — $C_{10}H_8N_2O_3S_2$, $M = 268.31$, monoclinic, space group $P2_1$, $a = 1.0018(2)$, $b = 0.9722(2)$, $c = 0.6262(1)$ nm, $\beta = 102.32(1)^\circ$, $V = 0.5958(2)$ nm³, $Z = 2$, $D_c = 1.496$ Mgm⁻³, $F(000) = 276$, $CuK\alpha$ radiation; $\lambda = 154.18$ pm, $\mu(CuK\alpha) = 3.998$ mm⁻¹.

Intensity Data, Structure Determination and Refinement. — Intensity data were collected on a Siemens AED diffractometer up to $\theta 70^\circ$ by the $\omega - 2\theta$ scan method (Ni-filtered $CuK\alpha$ radiation). 1262 reflections were measured, of which 1113, having intensities $I \geq 2\sigma(I)$ were used in the crystal analysis. The size of the crystal specimen was $0.07 \times 0.15 \times 0.49$ mm. No absorption correction was applied.

The structure was solved by direct methods using SHELX-76¹¹. An E-map computed by the use of phase set for 151 normalized structure factors having $E \geq 1.4$ revealed the positions of six atoms. A subsequent Fourier synthesis located the remaining non-hydrogen atoms. The refinement was carried out with isotropic ($R = 0.119$) and anisotropic thermal parameters, reducing R factor to 0.072. A difference Fourier synthesis revealed the positions of the hydrogen atoms, except H(5) and H(9). Including the hydrogen atoms with isotropic thermal parameters the refinement resulted in the final R factor of 0.051.

TABLE I

Fractional Atomic Coordinates ($\times 10^4$, for H Atoms $\times 10^3$) and Equivalent or Isotropic Thermal Parameters (pm^2) with Estimated Standard Deviations.

$$U_{eq} = 1/3 \sum_i \sum_j U_{ij} a_i^* a_j^* a_i \cdot a_j$$

	x	y	z	U_{eq}/U_{iso}
S(1)	2680(4)	560(4)	6902(4)	8.5(1)
S(2)	-648(2)	1986	-13(3)	4.6(1)
O(1)	3345(9)	2478(11)	4195(15)	11.6(4)
O(2)	-674(6)	1732(8)	-2278(9)	5.7(2)
O(3)	-625(7)	3378(6)	821(9)	5.6(2)
N(1)	703(8)	1244(8)	1353(11)	5.0(3)
N(2)	-6411(11)	-1319(14)	2596(16)	10.6(5)
C(1)	2520(12)	1724(12)	4669(16)	6.5(4)
C(2)	1091(10)	1348(10)	3710(13)	5.3(3)
C(3)	1119(13)	53(11)	5085(15)	6.8(4)
C(4)	-2065(8)	1148(9)	662(13)	4.6(3)
C(5)	-2622(11)	12(12)	-524(17)	6.6(4)
C(6)	-3719(10)	-642(12)	17(19)	7.2(4)
C(7)	-4288(9)	-172(10)	1703(15)	6.1(3)
C(8)	-3713(9)	962(11)	2907(16)	6.3(4)
C(9)	-2615(11)	1611(10)	2401(16)	6.0(4)
C(10)	-5447(10)	-828(14)	2236(18)	7.3(4)
H(1)	71(9)	44(6)	60(13)	8(2)
H(2)	59(7)	215(7)	427(12)	9(2)
H(31)	128(9)	-63(7)	387(14)	8(3)
H(32)	12(7)	12(9)	550(13)	14(3)
H(5)	-202(9)	-41(9)	-154(12)	10(3)
H(6)	-415(10)	-126(11)	-125(15)	10(3)
H(8)	-428(9)	133(9)	379(14)	13(3)
H(9)	-202(8)	234(7)	346(12)	12(3)

The positional parameters and equivalent or isotropic thermal parameters are given in Table I. The atomic scattering factors for non-hydrogen atoms were from ref. 12 and those for hydrogen atoms from ref. 13. Correction for the anomalous dispersion was taken from ref. 14. All calculations were carried out on the CDC Cyber 76 computer using the SHELX-76 program. Lists of structure factors and anisotropic thermal parameters are available from the authors on request.

DESCRIPTION AND DISCUSSION OF THE STRUCTURE

The molecular conformation and the atomic numbering scheme are shown in Figure 1. Interatomic distances and angles are given in Tables II and III, while selected torsion angles are listed in Table IV. Projection of the molecule along the bond $S(2)-N(1)$ is shown in Figure 2. Molecular packing with hydrogen bonding scheme is illustrated in Figure 3. Details on the least-square planes calculated for various parts of molecule are given in Table V.




Figure 1. Perspective view of the molecule.




Figure 2. Projection of the molecule looking along $S(2)-N(1)$ bond.




Figure 3. Molecular packing with hydrogen bonding scheme.

TABLE II
Interatomic Distances (pm)

S(1)—C(1)	177.8(9)	C(6)—C(7)	138.0(11)
S(1)—C(3)	180.4(10)	C(7)—C(8)	139.0(11)
C(1)—C(2)	147.5(12)	C(8)—C(9)	136.3(11)
C(1)—O(1)	118.9(10)	C(7)—C(10)	142.6(11)
C(2)—C(3)	152.2(10)	C(10)—N(2)	114.2(10)
N(1)—C(2)	144.8(8)	N(1)—H(1)	92(4)
N(1)—S(2)	160.9(6)	C(2)—H(2)	103(4)
S(2)—O(2)	143.4(4)	C(3)—H(31)	105(4)
S(2)—O(3)	144.9(5)	C(3)—H(32)	109(4)
S(2)—C(4)	176.5(7)	C(5)—H(5)	105(4)
C(4)—C(5)	138.0(10)	C(6)—H(6)	102(8)
C(4)—C(9)	139.6(9)	C(8)—H(8)	95(7)
C(5)—C(6)	137.3(12)	C(9)—H(9)	106(4)

The most important feature of this molecule is the conformation of the four-membered hetero ring. As expected the thietane ring is puckered. The puckering angle between planes S(1) C(1) C(2) and S(1) C(3) C(2) is 18.7(7) while that between planes C(1) C(2) C(3) and C(1) S(1) C(3) is 17.6(5)°. These angles in the present structure are significantly larger than those of the

TABLE III
Bond Angles (°)

C(1)—S(1)—C(3)	75.7(4)	O(3)—S(2)—C(4)	107.2(3)
S(1)—C(1)—C(2)	94.8(5)	N(1)—S(2)—C(4)	107.1(3)
S(1)—C(3)—C(2)	92.2(6)	S(2)—C(4)—C(5)	119.6(6)
C(1)—C(2)—C(3)	94.3(7)	S(2)—C(4)—C(9)	120.6(6)
O(1)—C(1)—S(1)	129.3(8)	C(9)—C(4)—C(5)	119.8(7)
O(1)—C(1)—C(2)	135.8(8)	C(4)—C(5)—C(6)	119.4(8)
C(1)—C(2)—N(1)	117.3(6)	C(5)—C(6)—C(7)	121.2(8)
C(3)—C(2)—N(1)	119.1(7)	C(6)—C(7)—C(8)	119.2(7)
C(2)—N(1)—S(2)	121.5(5)	C(7)—C(8)—C(9)	120.2(7)
O(2)—S(2)—O(3)	120.8(3)	C(8)—C(9)—C(4)	120.3(7)
O(2)—S(2)—N(1)	106.5(3)	C(6)—C(7)—C(10)	121.4(8)
O(2)—S(2)—N(1)	106.3(3)	C(8)—C(7)—C(10)	119.4(8)
O(2)—S(2)—C(4)	108.2(3)	C(7)—C(10)—N(2)	177.1(11)

TABLE IV
Selected Torsion Angles (°)

S(1)—C(1)—C(2)—C(3)	—13.9(8)
C(1)—C(2)—C(3)—S(1)	13.7(8)
C(1)—S(1)—C(3)—C(2)	—11.7(7)
C(3)—S(1)—C(1)—C(2)	12.1(7)
C(3)—S(1)—C(1)—O(1)	—165.8(14)
O(1)—C(1)—C(2)—C(3)	163.7(15)
S(1)—C(1)—C(2)—N(1)	—140.8(8)
O(1)—C(1)—C(2)—N(1)	37.6(19)
N(1)—C(2)—C(3)—S(1)	138.4(8)
C(1)—C(2)—N(1)—S(2)	—131.1(8)
C(3)—C(2)—N(1)—S(2)	116.2(9)
C(2)—N(1)—S(2)—C(4)	—67.5(8)
C(2)—N(1)—S(2)—O(2)	176.9(7)
C(2)—N(1)—S(2)—O(3)	46.9(8)
O(2)—S(2)—C(4)—C(5)	25.8(8)
O(2)—S(2)—C(4)—C(9)	—155.1(7)
O(3)—S(2)—C(4)—C(5)	157.7(7)
O(3)—S(2)—C(4)—C(9)	—23.2(8)
N(1)—S(2)—C(4)—C(5)	—88.6(8)
N(1)—S(2)—C(4)—C(9)	90.5(8)

corresponding derivatives with Cl, Br and H in *p*-position on the benzene ring¹⁻³ (13, 8.9 and 11.2°, respectively) but they are comparable with the values reported for some thietane derivatives with bulky substituents on vicinal carbons^{15,16} or for spiro compounds with thietane moiety¹⁷.

The values for S—C(sp³) and S—C(sp²) bond lengths are 180.4(10) and 177.8(9) pm, in agreement with the corresponding values reported for Br-derivative³ and those of Argay¹⁸ and Bocelli¹⁹. The C(1)—C(2) of 147.5(12) pm is much shorter than the C(2)—C(3) [152.2(10) pm]. The values of bond angles inside the ring show clearly the strain the ring is submitted to. The C(1)—S(1)—C(3) angle of 75.7(4)° is the smallest value reported so far for the thietane derivatives^{1-4,15-17,20-24}.

Atoms around the sulphur atom S(2) are arranged in a distorted tetrahedral coordination with enlarged O—S—O angle like in other sulphonamido compounds.

Projection along the S(2)—N(1) bond is illustrated in Figure 2. There is a great difference in orientation of the heterocyclic and benzene rings according to the analogous Cl and Br derivatives^{4,8}. These facts are also evident in the comparison of the corresponding torsion angles. The dihedral angle between the mean planes of two rings is 70.2(3)°.

The values of distances and angles in cyano group are in accordance with the known values in the literature²⁵⁻²⁷. The short C≡N distance is a typical problem in the X-ray structure analysis²⁸. This effect is due to thermal motion of the atoms and to the high electron density between the triple-bonded atoms. From Table V it is evident that these two atoms deviate significantly from the mean plane of the benzene ring.

TABLE V

Least-Squares Planes and Atomic Distances from These Planes. The Plane Equations Are of the Form $PX + QY + RZ = S$, Where X, Y and Z are in pm Relative to Orthogonal Axes

(a) Plane	P	Q	R	S
I S(1), C(1), C(2)	-44.5	71.0	54.6	190.4
II S(1), C(2), C(3)	-66.2	47.1	58.3	154.5
III S(1), C(1), C(3)	-62.2	63.1	46.4	120.2
IV C(1), C(2), C(3)	-46.7	53.0	70.7	202.2
V S(1), C(1), C(2), O(1)	43.4	-71.8	-54.4	-191.9
VI S(1), C(1), C(2), C(3), O(1)	51.4	-68.4	-51.7	-162.9
VII C(4), C(5), C(6), C(7), C(8), C(9)	-49.5	59.9	-63.0	147.6

(b) Deviations in pm From Planes:

V		VI		VII	
S(1)	0.0	S(1)	-1.4	C(4)	0.6
C(1)	-1.3	C(1)	-1.6	C(5)	-0.1
C(2)	0.3	C(2)	-13.3	C(6)	-1.0
O(1)	0.5	C(3)	21.4	C(7)	0.8
		O(1)	9.0	C(8)	-0.2
				C(9)	-0.7
				C(10)	3.1
				S(2)	0.6
				N(2)	10.8

(c) Dihedral Angle Between Planes (°)

I, II	18.7(7)	II, V	19.5(5)
III, IV	17.6(5)	VI, VII	70.2(3)

Molecules are connected in an infinitive chain along the *b* axis by hydrogen bonds N(1)—H(1)···O(3) of 309.5(9) pm; the angle is 172.(8)°. It is worth mentioning that the O atom participating in this hydrogen bond is from sulphonamido and not from thiolactone group, as in Br derivative³.

REFERENCES

1. I. Milinović, A. Bezjak, and D. Fleš, *Croat. Chem. Acta* **45** (1973) 551.
2. N. Galešić, M. Galešić, and E. Coffou, *Croat. Chem. Acta* **50** (1977) 283.
3. I. Matijašić, G. D. Andreotti, P. Sgarabotto, A. Bezjak, and D. Fleš, *Croat. Chem. Acta* **57** (1984) 621.

4. I. Matijašić, *Ph. D. Thesis*, PMF, Zagreb, 1984.
5. D. Fleš, A. Markovac-Prpić, and V. Tomašić, *J. Amer. Chem. Soc.* **80** (1958) 4654.
6. M. Dadić and D. Fleš, *Croat. Chem. Acta* **36** (1964) 37.
7. D. Fleš and V. Tomašić, *Polym. Lett.* **6** (1969) 809.
8. D. Fleš, V. Tomašić, M. Samsa, D. Ahmetović, B. Jerman, and M. Fleš, *J. Polym. Sci., Symposium No.* **42** (1973) 321.
9. D. Fleš, *Charged React. Polym.* **5** (*Opt. Act. Polym.*) (1979) 143.
10. Y. Chatani, T. Uchida, H. Tadokoro, K. Hayashi, M. Nishii, and S. Okamura, *J. Macromol. Sci. Phys.* **B2** (1968) 567.
11. G. M. Sheldrick, *SHELX-76. Program for Crystal Structure Determination*, Univ. of Cambridge, England, 1976.
12. D. T. Cromer and J. B. Mann, *Acta Cryst.* **A24** (1968) 321.
13. R. F. Stewart, E. R. Davidson, and W. T. Simpson, *J. Chem. Phys.* **42** (1965) 3175.
14. D. T. Cromer and D. Lieberman, *J. Chem. Phys.* **53** (1970) 1891.
15. V. Bertolasi and G. Gilli, *Acta Cryst.* **B34** (1978) 3403.
16. S. Kumakura and T. Kodama, *Bull. Chem. Soc. Jpn.* **48** (1975) 2339.
17. J. Guilhem, *Cryst. Struct. Comm.* **6** (1977) 93.
18. Gy. Argay, A. Kálmán, D. Lazar, and B. Ribár, *Acta Cryst.* **B33** (1977) 99.
19. G. Bocelli and M. F. Grenier-Loustalot, *Cryst. Struct. Comm.* **11** (1982) 85.
20. H. Nakai and H. Koyama, *J. Chem. Soc. B* (1971) 1525.
21. H. Nakai and H. Koyama, *J. Chem. Soc. Perkin II* (1972) 248.
22. C. D. Shirell and D. E. Williams, *Acta Cryst.* **B29** (1973) 2128.
23. E. Schaumann, H. G. Bäuch, and G. Adiwidjaja, *Angew. Chem. Int. Ed. Engl.* **20** (1981) 613.
24. K. H. Jogun and J. J. Stezowski, *Acta Cryst.* **B35** (1979) 2310.
25. Nguyen-Huy-Dung and J. Etienne, *Acta Cryst.* **B34** (1978) 683.
26. R. Schlodder, J. A. Ibers, M. Lenarda, and M. Graziani, *J. Amer. Chem. Soc.* **96** (1974) 6893.
27. T. Higashi and K. Osaki, *Acta Cryst.* **B33** (1977) 607.
28. R. A. Loghry and S. H. Simonsen, *Acta Cryst.* **B32** (1976) 1505.

SAŽETAK

Kristalna i molekularna struktura (S)- α -(*p*-cijanobenzensulfonamido)- β -propiotiolaktona

I. Matijašić, G. D. Andreotti, P. Sgarabotto, A. Bezjak i D. Fleš

Spoj kristalizira u monoklinskom sustavu sa $a = 1,0018(2)$, $b = 0,9722(2)$, $c = 0,6262$ pm i $\beta = 102,32(1)^\circ$, $Z = 2$ u prostornoj grupi $P2_1$. Struktura je riješena direktnim metodama upotrebom 1113 refleksa sa $I \geq 2\sigma(I)$ i utočnjena do $R = 0,051$. Heterociklički četveročlani prsten ima kut izvijanja od $18,7(7)^\circ$. Diedarski kut između ravnina položenih β -propiotiolaktonskim i benzenskim prstenom iznosi $70,2(3)^\circ$. Struktura se sastoji od molekula povezanih duž osi b intermolekulskom vodikovom vezom N—H...O od 309,5(9) pm.